

Linear Programming I

Why Linear Programming rocks:

- Incredibly general: Every problem we've seen so far can be formulated as a linear program.
- Computational tractable
 - In theory: Can be solved in polynomial time
 - In practice: Fast with input sizes up into the millions!
- Contains many properties that can be turned into useful algorithmic paradigms and analysis:
 - Duality:
 - * Solve an easier equivalent problem.
 - * How do we know when we're done?
 - Complementary Slackness and Strong Duality: something is optimal!

How to Think About Linear Programming

Comparison to Systems of Linear Equations

Remember systems of equations, with a_{ij} 's and b_i 's given. We used Gaussian elimination; to find feasible x_1, \dots, x_n , or correctly reports that no feasible solution exists.

$$\begin{aligned}a_{11}x_1 + a_{12}x_2 + \dots + a_{1n}x_n &= b_1 \\a_{21}x_1 + a_{22}x_2 + \dots + a_{2n}x_n &= b_2 \\&\vdots \\a_{m1}x_1 + a_{m2}x_2 + \dots + a_{mn}x_n &= b_m.\end{aligned}$$

What about linear *inequalities*? The point of linear programming is to solve these, and when there are multiple feasible solutions, we would like to compute the “best” one.

Ingredients of a Linear Program

Ingredients of a Linear Program

- a. *Decision variables* $x_1, \dots, x_n \in \mathbb{R}$.

b. *Linear constraints*, each of the form

$$\sum_{j=1}^n a_j x_j \quad (*) \quad b_i,$$

where $(*)$ could be \leq , \geq , or $=$.

c. A *linear objective function* of the form

$$\max \sum_{j=1}^n c_j x_j \quad \text{or} \quad \min \sum_{j=1}^n c_j x_j.$$

Comments:

- The a_{ij} 's, b_i 's, and c_j 's are
- The x_j 's are
- On “ \leq ” and “ \geq ” inequalities
- On equality constraints
- On “min” and “max” cases for the objective function

What's not allowed in a linear program? Non-linear terms like x_j^2 , $x_j x_k$, $\log(1+x_j)$, etc. We're only allowed to have x_j multiplied by a constant.

A Simple Example

To make linear programs more concrete and develop your geometric intuition about them, let's look at a toy example. (Many “real” examples of linear programs are coming shortly.) Suppose there are two decision variables x_1 and x_2 —so we can visualize solutions as points (x_1, x_2) in the plane. See Figure ???. Let's consider the (linear) objective function of maximizing the sum of the decision variables:

$$\max x_1 + x_2.$$

We'll look at four (linear) constraints:

$$\begin{aligned}x_1 &\geq 0 \\x_2 &\geq 0 \\2x_1 + x_2 &\leq 1 \\x_1 + 2x_2 &\leq 1.\end{aligned}$$




Figure 1: A toy example of a linear program.

The first two inequalities restrict feasible solutions to the non-negative quadrant of the plane. The second two inequalities further restrict feasible solutions to lie in the shaded region depicted in Figure ???. Geometrically, the objective function asks for the feasible point furthest in the direction of the coefficient vector $(1, 1)$ —the “most northeastern” feasible point. Eyeballing the feasible region, this point is $(\frac{1}{3}, \frac{1}{3})$, for an optimal objective function value of $\frac{2}{3}$. This is the “last point of intersection” between a level set of the objective function and the feasible region (as one sweeps from southwest to northeast).

Geometric Intuition

While it’s always dangerous to extrapolate from two or three dimensions to an arbitrary number, the geometric intuition above remains valid for general linear programs, with an arbitrary number of dimensions (i.e., decision variables) and constraints. Even though we can’t draw pictures when there are many dimensions, the relevant algebra carries over without any difficulties. Specifically:

- a. A linear constraint in n dimensions corresponds to a halfspace in \mathbb{R}^n . Thus a feasible region is an intersection of halfspaces, the higher-dimensional analog of a polygon.¹
- b. When there is a unique optimal solution, it is a _____ of the feasible region.

A few edge cases:

- a. There might be no feasible solutions at all. For example, if we add the constraint $x_1 + x_2 \geq 1$ to our toy example, then there are no longer any feasible solutions. Linear programming algorithms correctly detect when this case occurs.
- b. The optimal objective function value is unbounded ($+\infty$ for a maximization problem, $-\infty$ for a minimization problem). Note a necessary but not sufficient condition for this case is that the feasible region is unbounded. For example, if we dropped the constraints $2x_1 + x_2 \leq 1$

¹A finite intersection of halfspaces is also called a “polyhedron;” in the common special case where the feasible region is bounded, it is called a “polytope.”

and $x_1 + 2x_2 \leq 1$ from our toy example, then it would have unbounded objective function value. Again, linear programming algorithms correctly detect when this case occurs.

- c. The optimal solution need not be unique, as a “side” of the feasible region might be parallel to the levels sets of the objective function. Whenever the feasible region is bounded, however, there always exists an optimal solution that is a vertex of the feasible region.²

Examples

Example 1: Grain Nutrients

Suppose BU has hired you to optimize nutrition for campus dining. There are two possible grains they can offer, grain 1 and grain 2, and each contains the macronutrients found in the table below, plus cost per kg for each of the grains.

| Macros | Starch | Proteins | Vitamins | Cost (\$/kg) |
|---------|--------|----------|----------|--------------|
| Grain 1 | 5 | 4 | 2 | 0.6 |
| Grain 2 | 7 | 2 | 1 | 0.35 |

The nutrition requirement per day of starch, proteins, and vitamins is 8, 15, and 3 respectively. Determine how much of each grain to buy such that BU spends as little but meets its nutrition requirements.

Decision variables:

Objective:

Constraints:

²There are some annoying edge cases for unbounded feasible regions, for example the linear program $\max(x_1 + x_2)$ subject to $x_1 + x_2 = 1$.

Example 2: Transportation

You're working for a company that's producing widgets among two different factories and selling them from three different centers. Each month, widgets need to be transported from the factories to the centers. Below are the transportation costs from each factory to each center, along with the monthly supply and demand for each factory and center respectively. Determine how to route the widgets in a way that minimizes transportation costs.

| Transit Cost | Center 1 | Center 2 | Center 3 |
|--------------|----------|----------|----------|
| Factory 1 | 5 | 5 | 3 |
| Factory 2 | 6 | 4 | 1 |

- The supply per factory is 6 and 9 respectively.
- The demand per center is 8, 5, and 2 respectively.

Decision variables:

Objective:

Constraints:

Converting to Normal Form

The “Normal Form” of a Linear Program looks like:

$$\begin{aligned} \max \quad & \mathbf{c}^T \mathbf{x} \\ \text{subject to} \quad & \mathbf{Ax} \leq \mathbf{b} \end{aligned}$$

Our Transportation problem had the LP:

$$\begin{aligned} \min \quad & 5x_{11} + 5x_{12} + 3x_{13} + 6x_{21} + 4x_{22} + 1x_{23} \\ \text{subject to} \quad & x_{11} + x_{12} + x_{13} = 6 && \text{(Factor 1 supply)} \\ & x_{21} + x_{22} + x_{23} = 9 && \text{(Factor 2 supply)} \\ & x_{11} + x_{21} = 8 && \text{(Center 1 demand)} \\ & x_{12} + x_{22} = 5 && \text{(Center 2 demand)} \\ & x_{13} + x_{23} = 2 && \text{(Center 3 demand)} \\ & x_{ij} \geq 0 && \text{(non-negativity)} \end{aligned}$$

How can we convert it to normal form—a maximization problem with all less-than-or-equal-to constraints?

First observe that $x_{11} + x_{12} + x_{13} = 6$ is equivalent to having both inequalities

But, we need both to be \leq inequalities! We transform them to

The resulting LP in normal form is:

Writing Problems We Know as Linear Programs

Independent Set

Given a graph $G = (V, E)$, each vertex i has weight w_i , find a maximum weighted *independent set*.
 S is an independent set if it does not contain both i and j for $(i, j) \in E$.

a. Decision variables: What are we try to solve for?

b. Constraints:

c. Objective function:

The linear program:

Integer programs vs. linear relaxations: