

Prophet Inequalities

You're at a casino gambling, and are offered to play the following game. Items will arrive one-by-one. As an item arrives, you see its value. You may only take a single item, and once you take an item, the game ends. A priori, you know the *distribution* of each item. At some point there will be a red item with the red distribution of values, and at some point there will be a blue item with the blue distribution of values, and so forth. However, you do not know the order of items (it is adversarial), and you do not know the exact values of the items (they are drawn from their specific distributions). Your goal is to come up with an algorithm that competes with the *prophet* who is all knowing, so knows the realization of values and the arrival order.

That is, n items will arrive in adversarial order. Item i (which is a label, not necessarily the order) has value v_i drawn from known distribution F_i . Your goal is to determine an algorithm ALG such that the value you get from gambling competes with the prophet who always gets $\max_i v_i$. However, your competition is over the randomness of the values that are drawn, so you only have to compete with $\text{OPT} = \mathbb{E}_{\mathbf{v}}[\max_i v_i]$.




Figure 1: The prophet inequality problem.

To summarize:

- Goal: Pick one item; maximize its value.
- Gambler knows distribution for each item.
- Order is adversarial.
- Inspect each item online (see v_i) and irrevocably decide whether to take or pass forever.
- Compete with $\text{OPT} = \mathbb{E}_{\mathbf{v}}[\max_i v_i]$.

The Prophet Inequality problem was posed by Samuel-Cahn '84 [7], with the original solution and analysis that we'll see by Krengel Sucheston '78 [6] and Garling.

Prove the following.

Theorem 1. *There is a threshold algorithm ALG such that when the gambler takes an item if and only if its value is above T , $\text{ALG} \geq \frac{1}{2}\text{OPT}$.*

Determine what threshold T to use and prove this statement using the following steps:

1. Divide what the algorithm yields from an item (in expectation) into exactly the threshold and the surplus above the threshold.
2. Lower bound your surplus term.
3. Set your threshold in order to combine like-terms and have OPT pop out.

Note: Can you find two different thresholds that give this same approximation?

Proof. We consider two different ways to set the threshold. Let p denote the probability that *some* (at least one) $v_i \geq T$ for $i \in [n]$.

We reinterpret our problem as follows: we have a single item to sell, and n buyers arrive online with their values for the item drawn from distributions. We price the item at T . Observe that the first term above is the *expected revenue* earned from selling the item to the buyers at T , and the second term is the *expected utility*, or surplus. Welfare is equal to revenue + utility.

Exercise: We can see this as a mechanism for a buyer to maximize social welfare [3]. Could you design a mechanism to maximize revenue using the prophet inequality?

[Hint: Use virtual values.]

Other Extensions: Suppose the gambler can take k items, or a matroid of items [4, 5]. What algorithm should they use to determine which items to take as they arrive online?

The Multidimensional Extension [1, 2]

Imagine now that a seller has m goods to sell, and n buyers will arrive online one at a time. They have a combinatorial value function $v_i : 2^m \rightarrow \mathbb{R}_{\geq 0}$ and each function v_i is drawn from a known

prior distribution. Our goal is to set prices such that the welfare of the allocation is maximized.

Notationally:

- For the allocation \mathbf{x} such that x_i is the allocation to bidder i , set a pricing function $p : 2^m \rightarrow \mathbb{R}_{\geq 0}$ such that under online arrival, the buyer's allocations maximize $\sum_i v_i(x_i)$.
- Let \mathbf{x} denote a partial allocation rule, for instance, the allocation of items after arrival so far.
- Let \mathbf{x}' denote a second allocation rule that, combined with the partial allocation rule of \mathbf{x} , is still feasible. That is, in this online context, \mathbf{x}' is a feasible allocation of the remaining items not allocated by \mathbf{x} .
- Let $\text{OPT}(\mathbf{v} \mid \mathbf{x})$ denote the optimal welfare of all allocations that are feasible with \mathbf{x} , i.e., the value remaining (not including the value from \mathbf{x}) when \mathbf{x} is already committed to. Formally,

$$\text{OPT}(\mathbf{v} \mid \mathbf{x}) = \max_{\mathbf{x}' : \mathbf{x} \cup \mathbf{x}' \text{ feas.}} v_i(x'_i).$$

Definition 1. A pricing rule p is (α, β) -balanced with respect to valuation profile $\mathbf{v} = (v_1, v_2, \dots, v_n)$ if, for all feasible allocations \mathbf{x} and \mathbf{x}' such that $\mathbf{x} \cup \mathbf{x}'$ is feasible:

1. $\sum_i p(x_i) \geq \frac{1}{\alpha} \cdot \underbrace{(\text{OPT}(\mathbf{v}) - \text{OPT}(\mathbf{v} \mid \mathbf{x}))}_{\text{value lost due to allocating } \mathbf{x}}.$
2. $\sum_i p(x'_i) \leq \beta \cdot \underbrace{\text{OPT}(\mathbf{v} \mid \mathbf{x})}_{\text{value remaining after allocating } \mathbf{x}}.$

Prices are *weakly balanced* if the second condition is relaxed to $\text{OPT}(\mathbf{v})$.

Theorem 2. If a pricing rule p is (α, β) -balanced with respect to valuations \mathbf{v} , then posting prices $\delta \cdot p$ guarantees value at least $\frac{1}{\alpha\beta+1} \cdot \text{OPT}(\mathbf{v})$ for $\delta = \frac{\alpha}{\alpha\beta+1}$.

Sanity check: In the single-item case, a price $p = \max_i v_i$ is $(1, 1)$ -balanced. Which implies?

Proof. Let \mathbf{x} be the allocation sold, and let \mathbf{x}' be the allocation achieving $\text{OPT}(\mathbf{v} \mid \mathbf{x})$. Then we get that

$$\text{REVENUE} =$$

$$\text{UTILITY} \geq$$

□

Extension to unknown values: For (α, β) -balanced p^v , let $p = \mathbb{E}_{v \sim F}[p^v]$. Then posting prices δp guarantees expected value $\frac{1}{\alpha\beta+1} \cdot \mathbb{E}[\text{OPT}(\mathbf{v})]$. (For weakly balanced, it's $\frac{1}{4\alpha\beta}$.)

Construction of Balanced Prices: If any individual's allocation is bounded at k items, and if \mathbf{x}^* is the optimal allocation, then we set

$$p_j = \begin{cases} \frac{1}{|x_i^*|} v_i(x_i^*) & j \in x_i^* \\ 0 & \text{otherwise} \end{cases}.$$

Claim 1. These are weakly $(k, 1)$ -balanced.

Proof.

References

- [1] Paul Dütting, Michal Feldman, Thomas Kesselheim, and Brendan Lucier. Prophet inequalities made easy: Stochastic optimization by pricing non-stochastic inputs. In Chris Umans, editor, *58th IEEE Annual Symposium on Foundations of Computer Science, FOCS 2017, Berkeley, CA, USA, October 15-17, 2017*, pages 540–551. IEEE Computer Society, 2017.
- [2] Michal Feldman, Nick Gravin, and Brendan Lucier. Combinatorial auctions via posted prices. In *Proceedings of the Twenty-Sixth Annual ACM-SIAM Symposium on Discrete Algorithms*, pages 123–135. SIAM, 2015.
- [3] Mohammad Taghi Hajiaghayi, Robert Kleinberg, and Tuomas Sandholm. Automated online mechanism design and prophet inequalities. In *Proceedings of the 22nd National Conference on Artificial Intelligence - Volume 1, AAAI'07*, page 58?65. AAAI Press, 2007.
- [4] Robert Kleinberg and S. Matthew Weinberg. Matroid prophet inequalities. In Howard J. Karloff and Toniann Pitassi, editors, *Proceedings of the 44th Symposium on Theory of Computing Conference, STOC 2012, New York, NY, USA, May 19 - 22, 2012*, pages 123–136. ACM, 2012.

- [5] Robert Kleinberg and S. Matthew Weinberg. Matroid prophet inequalities and applications to multi-dimensional mechanism design. *Games Econ. Behav.*, 113:97–115, 2019.
- [6] Ulrich Krengel and Louis Sucheston. Semiamarts and finite values. *Bulletin of the American Mathematical Society*, 83(4):745–747, 1977.
- [7] Ester Samuel-Cahn. Comparison of threshold stop rules and maximum for independent non-negative random variables. *The Annals of Probability*, pages 1213–1216, 1984.